

This is a repository copy of *Delirium detection in older acute medical inpatients : a 2 multicentre prospective comparative diagnostic test accuracy 3 study of the 4AT and the Confusion Assessment Method*.

White Rose Research Online URL for this paper:

<https://eprints.whiterose.ac.uk/147621/>

Version: Accepted Version

---

**Article:**

Shenkin, Susan D, Fox, Christopher, Godfrey, Mary et al. (14 more authors) (2019) Delirium detection in older acute medical inpatients : a 2 multicentre prospective comparative diagnostic test accuracy 3 study of the 4AT and the Confusion Assessment Method. BMC Medicine. ISSN 1741-7015

---

**Reuse**

Items deposited in White Rose Research Online are protected by copyright, with all rights reserved unless indicated otherwise. They may be downloaded and/or printed for private study, or other acts as permitted by national copyright laws. The publisher or other rights holders may allow further reproduction and re-use of the full text version. This is indicated by the licence information on the White Rose Research Online record for the item.

**Takedown**

If you consider content in White Rose Research Online to be in breach of UK law, please notify us by emailing [eprints@whiterose.ac.uk](mailto:eprints@whiterose.ac.uk) including the URL of the record and the reason for the withdrawal request.

# **Title: Delirium detection in older acute medical inpatients: a multicentre prospective comparative diagnostic test accuracy study of the 4AT and the Confusion Assessment Method**

Susan D Shenkin (Susan.Shenkin@ed.ac.uk)<sup>1</sup>

Christopher Fox (Chris.Fox@uea.ac.uk)<sup>2</sup>

Mary Godfrey (m.godfrey@leeds.ac.uk)<sup>3</sup>

Najma Siddiqi (najma.siddiqi@york.ac.uk)<sup>4</sup>

Steve Goodacre (s.goodacre@sheffield.ac.uk)<sup>5</sup>

John Young (John.Young@bthft.nhs.uk)<sup>6</sup>

Atul Anand (atul.anand@ed.ac.uk)<sup>7</sup>

Alasdair Gray (alasdaigray@blueyonder.co.uk)<sup>8</sup>

Janet Hanley (janet.hanley@nhs.net)<sup>9</sup>

Allan MacRaid (Allan.MacRaid@nhslothian.scot.nhs.uk)<sup>8</sup>

Jill Steven (Jill.Steven@nhslothian.scot.nhs.uk)<sup>8</sup>

Polly L Black (Polly.Black@nhslothian.scot.nhs.uk)<sup>8</sup>

Zoë Tieges (Zoe.Tieges@ed.ac.uk)<sup>1</sup>

Julia Boyd (Julia.Boyd@ed.ac.uk)<sup>10</sup>

Jacqueline Stephen (Jacqueline.Stephen@ed.ac.uk)<sup>10</sup>

Christopher J Weir (Christopher.Weir@ed.ac.uk)<sup>10</sup>

Alasdair MJ MacLulich (a.maclulich@ed.ac.uk)<sup>1</sup>

## **Institutional addresses**

1. Geriatric Medicine, University of Edinburgh, Edinburgh, UK
2. Norwich Medical School, University of East Anglia, Norfolk, UK
3. Elderly Care and Rehabilitation, and, Institute of Health Sciences, University of Leeds, Leeds, UK
4. Department of Health Sciences, University of York, York, Hull York Medical School, York and Bradford District Care NHS Foundation Trust, Bradford, UK
5. School of Health and Related Research (ScHARR), University of Sheffield, Sheffield, UK
6. Academic Unit of Elderly Care and Rehabilitation, University of Leeds, Leeds, UK
7. Cardiovascular Sciences and Geriatric Medicine, University of Edinburgh, Edinburgh, UK
8. Emergency Medicine Research Group (EMERGE), NHS Lothian, Edinburgh, UK
9. Health and Social Care, Edinburgh Napier University, Edinburgh, UK
10. Edinburgh Clinical Trials Unit, Usher Institute of Population Health Sciences and Informatics, University of Edinburgh, Edinburgh, UK

**Corresponding Author**

Professor Alasdair MJ MacLulich

Professor of Geriatric Medicine, University of Edinburgh

Honorary Consultant in Geriatric Medicine, Royal Infirmary of Edinburgh

Edinburgh Delirium Research Group

Geriatric Medicine Unit, University of Edinburgh

Room S1642, Royal Infirmary of Edinburgh

51, Little France Crescent, Edinburgh EH16 4SA

Tel: 0131 242 6481

Email: [a.maclulich@ed.ac.uk](mailto:a.maclulich@ed.ac.uk)

# Abstract

## Background

Delirium affects >15% of hospitalised patients but is grossly underdetected, contributing to poor care. The 4 'A's Test (4AT; [www.the4AT.com](http://www.the4AT.com)) is a short delirium assessment tool designed for routine use without special training. The primary objective was to assess the accuracy of the 4AT for delirium detection. The secondary objective was to compare the 4AT with another commonly-used delirium assessment tool, the Confusion Assessment Method (CAM).

## Methods

This was a prospective diagnostic test accuracy study set in Emergency Departments or acute medical wards involving acute medical patients aged  $\geq 70$ . All those without acutely life-threatening illness or coma were eligible. Patients underwent (1) reference standard delirium assessment based on DSM-IV criteria and (2) were randomised to either the index test (4AT, scores 0-12; prespecified score of  $>3$  considered positive) or the comparator (CAM; scored positive or negative), in a random order, using computer-generated pseudo-random numbers, stratified by study site, with block allocation. Reference standard and 4AT or CAM assessments were performed by pairs of independent raters blinded to the results of the other assessment.

## Results

843 individuals were randomised: 21 withdrew, 3 lost contact, 32 indeterminate diagnosis, 2 missing outcome; 785 were included in the analysis. Mean age was 81.4 (SD 6.4) years.

12.1% (95/785) had delirium by reference standard assessment, 14.3% (56/392) by 4AT, and 4.7% (18/384) by CAM. The 4AT had an area under the receiver operating characteristic curve of 0.90 (95% CI 0.84-0.96). The 4AT had a sensitivity of 76% (95% CI 61-87%) and a specificity of 94% (95% CI 92-97%). The CAM had a sensitivity of 40% (95% CI 26-57%) and a specificity of 100% (95% CI 98-100%).

## Conclusions

The 4AT is a short, pragmatic tool which can help improving detection rates of delirium in routine clinical care.

## Registration

International standard randomised controlled trial number (ISRCTN) 53388093.

## Funding source

National Institute of Health Research Health Technology Assessment Programme (NIHR HTA) grant number 11/143/01. CJW was also supported in this work by NHS Lothian via Edinburgh Clinical Trials Unit. The views expressed are those of the authors and not necessarily those of the NHS, the NIHR or the Department of Health and Social Care.

## Keywords

Delirium; diagnostic test accuracy; 4AT; Confusion Assessment Method (CAM); sensitivity; specificity; hospital

# Background

Delirium is a severe neuropsychiatric syndrome, usually triggered by underlying medical illness, surgery or drugs, which affects at least 15% of hospital inpatients [1-4]. It is more common in older people [5] and people with dementia [6]. Delirium comprises acute onset of disturbances in arousal, attention and other domains of cognition, hallucinations and delusions [7, 8]. Delirium is important because as well as being highly prevalent in hospitalised patients, it strongly predicts poor outcomes such as falls, other medical complications, new institutionalisation and mortality [1, 6, 9-13]. It is also associated with patient and carer distress [14-16]. At least two-thirds of cases are not identified in Emergency Department and general medical settings [17-21]. The reasons for this include time constraints, and lack of education and training [22-24]. Because formal psychiatric assessment for delirium diagnosis takes considerable time, guidelines and pathways advocate use of brief assessment tools for delirium detection. Two assessment tools extensively used in clinical practice are the 4 'A's Test (4AT) and the short form of the Confusion Assessment Method (CAM).

The 4AT [25, 26] comprises four items: (A) Alertness, (B) Abbreviated Mental Test-4; (C) Attention (Months Backwards test); and (D) Acute change or fluctuating course [25, 27], Figure 1. The 4AT was not derived directly from a single set of diagnostic criteria; rather it has items that inform the core features of standard diagnostic criteria. It has a score range of 0-12, with scores of 4 or more ( $>3$ ) suggesting possible delirium. The structure of the 4AT is designed such that there are different ways of reaching an overall positive score ( $>3$ ). Items (A) and (D) each give a score of 0 if negative, and 4 if positive. The rationale for items (A) and (D) individually potentially triggering an overall positive 4AT score is that altered

arousal and acute change are both highly specific features of delirium [28-30]. The AMT-4 (B) gives a score of 1 for one mistake, and 2 for two or more mistakes or if the patient is untestable. The attention test (C) gives a score of 1 if unable to complete 7 months backwards, and 2 if untestable. Therefore patients who perform poorly or are untestable on both cognitive tests (B+C), score 4 from items (B) + (C), triggering further assessment for delirium. The rationale for the (B) and (C) scoring is that many patients with delirium are unable to undergo cognitive testing because of reduced arousal or other reasons [31, 32], and they would be unscorable or scored as negative on assessments that require cognitive testing, but the 4AT identifies that further assessment is required. The 4AT takes around 2 minutes and does not require special training. It is recommended in several pathways and guidelines and is in wide routine clinical use in the UK and internationally. Since publication on a dedicated website [26] in 2011, the 4AT has to date been evaluated in eight validation studies [25, 33-39] involving a total of 2577 patients, 479 with delirium. These studies have used varying designs, reference standards, clinical populations, and inclusion criteria. Sensitivities are reported as 83-100% and specificities ranging from 70-99%.

#### **Figure 1. The 4 'A's Test (4AT)**

The CAM (short form) [28] comprises brief cognitive testing and interview followed by a four-item algorithm in which four DSM-III-R criteria for delirium are rated as being present or absent: (A) acute onset and fluctuating course, (B) inattention, (C) disorganised thinking and (D) level of consciousness. To score positive on the CAM, both (A) and (B) must be positive, plus either or both of (C) and (D). The CAM requires specific training in rating



each of the features. The cognitive testing which is carried out before completing the algorithm is not specified in the manual [40]. With the pre-algorithm interview and cognitive testing it takes 5-10 minutes to complete [40]. The CAM is included in multiple international guidelines and pathways, including the UK NICE Guidelines on Delirium published in 2010 [41]. According to published systematic reviews [42-46] and a literature review carried out on 19 Feb 2019, the CAM has been evaluated in 22 validation studies since publication in 1990 [31, 47-67], with a total of 2437 participants (620 with delirium). As with the 4AT validation studies, these studies vary in design, population, etc. The reported range of sensitivities for delirium detection is 13-100%, and the range of specificities 84-100%.

The primary objective of the present study was to conduct an evaluation of the diagnostic accuracy of the 4AT for delirium against a reference standard based on DSM-IV in patients aged 70 years and over recently admitted to hospital recruited prospectively. The secondary objective was to compare the diagnostic test accuracy of the 4AT and CAM. The rationale for performing the comparison is that the 4AT and CAM are both widely used and recommended, yet the 4AT and CAM differ in their scoring systems, and the 4AT offers potential advantages include a shorter testing duration and no need for specific training, and a process for handling untestable patients. Given these differences, it is of interest to practitioners and researchers to know if performance of the 4AT is at least equivalent to the CAM. Additionally, both the 4AT and the CAM have been evaluated in multiple validation studies, but there are no published studies comparing performance of these tools under the same study conditions.

## Methods

We followed the Standards for Reporting Diagnostic Accuracy (STARD) 2015 guidelines [68] for reporting diagnostic accuracy studies. The study was registered: International standard randomised controlled trial number (ISRCTN) 53388093. UK Clinical Research Network ID: 19502, and the protocol published before database lock and statistical analysis [69]. The objectives described in the protocol not reported here (e.g. 12 week outcomes) will be disseminated separately.

### Study design: overview

The study protocol has been published [69]. In summary, patients aged 70 or over in Emergency Departments or acute general medical wards were prospectively recruited in three UK sites (Edinburgh, Bradford, and Sheffield). Each patient underwent (a) a reference standard delirium assessment lasting up to 20 minutes, and (b) either the 4AT or the CAM. Participants were randomised to the 4AT or the CAM and also to the ordering of the reference standard and the 4AT or CAM assessment. The study flowchart is shown in Figure 2.

### Figure 2. Diagnostic accuracy study: overview flowchart

## Participants

Potentially eligible participants were those without acutely life-threatening illness or coma, in the Emergency Department or acute general medical wards. Initially the recruitment windows were four hours for the Emergency Department and 24 hours for the acute general medical wards. Four months after study commencement (from 19 February 2016) these were extended to 12 hours and 96 hours respectively to facilitate recruitment, particularly with respect to seeking proxy consent. The potential impact of this was explored in planned subgroup analyses.

Patients were recruited by researchers between 0800 and 2200, Monday to Friday, from eligible patients identified by the clinical team. Patients were initially approached alphabetically, then in approximately the last third of the recruitment period, through liaison with clinical staff, prioritising those at higher risk of delirium on clinical grounds (e.g. older age, likely to be admitted, higher degree of ongoing acute and chronic illnesses) to obtain a more representative sample of participants [69] because interim analysis found a lower than expected rate of recruitment of patients who lacked capacity and thus required proxy consent. These changes to the recruitment processes were approved by the Trial Steering Committee. Informed consent was sought by trained researchers. Where the potential participant lacked capacity to consent, recruitment proceeded under the provisions of the Mental Capacity Act, 2005 in England or Adults with Incapacity (Scotland) Act, 2000, using an appropriate personal or nominated consultee, guardian, welfare attorney or nearest relative [69].

## Test methods

Researchers were nurses or trained graduate clinical research associates who underwent a systematic and detailed training process involving teaching on delirium and dementia assessment [69]. Additionally, training on the CAM was provided according to the guidance given in the CAM instruction manual [40]. Specific training on the 4AT was not provided as the tool was designed such that this is not required. The study team had regular teleconferences to discuss the conduct of the study.

The reference standard assessment was based on DSM-IV. These diagnostic criteria were used rather than DSM-5 because the study, ethics applications and training procedures were initiated at a similar time to publication of DSM-5 and it was not yet in use by the study team; because DSM-IV had been used in large numbers of delirium studies thus providing more direct comparability with the existing literature; and because of concern that there was insufficient time to develop and test valid methods for reference standard assessment using DSM-5. The reference standard drew from several sources of information including all items from the Delirium Rating Scale-Revised-98 (DRS-R98)[70] and using the instructions from the manual, which include raters seeking informant history and inspection of clinical records, and a set of neuropsychological tests designed to detect core features of delirium [71, 72] comprising Observational Scale for Level of Arousal [29, 73], the Richmond Agitation-Sedation Scale [74], Digit Span [75], the Vigilance A test [76], the DelApp objective test of attention [77-79], and standard object naming and orientation questions. These assessments were used together to inform a binary diagnosis of delirium based on DSM-IV criteria. The initial diagnosis was recorded by the researcher performing the assessment for the purposes of providing immediate information to the clinical team. These initial results of the reference standard assessment were provided by this researcher to the

1 clinical teams after the study 4AT or CAM were completed, through both an entry in the  
2 clinical notes, and a verbal discussion. The final and definitive ascertainment for the study  
3 was performed later, via expert consensus from a panel comprising ZT (a psychologist),  
4 SDS (a geriatrician) and AMJM (a geriatrician), each with many years of experience of  
5 delirium assessment (>1000 episodes individually). This final ascertainment was based on  
6 the information generated by the reference standard assessment in relation to the DSM-IV  
7 criteria, blinded to knowledge of whether the participant had undergone the 4AT or CAM,  
8 or the results of these tests. Where there was disagreement, the panel discussed each case  
9 using the available information and reached consensus. Where the reference standard  
10 assessment results did not provide enough information to provide a definite diagnosis of  
11 delirium, the ascertainment was judged to be indeterminate.

12 The 4AT was scored according to the guidance notes on the 4AT form [26], using a  
13 combination of sources of available information including casenotes, informants, and  
14 bedside assessment. As per the initial design of the 4AT, scores of >3 were used to indicate  
15 possible delirium. If patients were unable to undergo cognitive testing because of reduced  
16 arousal, the cognitive items 2 and 3 were scored as 'untestable' and each given a score of 2  
17 as per the 4AT guidance notes.

18 The CAM algorithm was scored following an interview and set of cognitive tests, and other  
19 sources of available information including casenotes and informants, as recommended in  
20 the CAM instruction manual [40]. The interview comprised general questions about the  
21 patient's hospital stay followed by a set of cognitive tests comprising: days of the week  
22 backwards, counting from 20 down to 1, orientation (current day, identifying if it is day or  
23 night, current year, last meal, how long in hospital, city, name of the hospital, floor of the  
24 hospital), memory (3 word recall immediately, up to 3 trials until all 3 words recalled or 3

1 trials repeated; then recall at 5 minutes), and clock drawing. The CAM algorithm was scored  
2 as per the instruction manual. Where an item could not be assessed, for example, if the  
3 patient was unable to speak or write and thus could not undergo assessment for disorganised  
4 thinking (see instruction manual), the item was scored as negative.

5 The presence of dementia was sought through either a formal diagnosis of dementia in the  
6 clinical records and/or, when possible, the Informant Questionnaire on Cognitive Decline  
7 in the Elderly (IQCODE) using a cut-off score of  $\geq 3.44$  [80].

## 8 **Ordering of reference standard delirium assessment, 4AT and CAM**

9 After the consent process was complete, participants were randomised in a 1:1 ratio to (a)  
10 reference standard first then either 4AT or CAM or (b) either 4AT or CAM first then  
11 reference standard via a secure online system using computer-generated pseudo-random  
12 numbers, stratified by study site, with block allocation. The reference standard assessment  
13 was performed by the researcher who conducted the capacity assessment and consenting  
14 process. A different researcher from the one performing the reference standard assessment  
15 performed either the 4AT or the CAM. Researchers performed the 4AT or the CAM  
16 according to the randomisation, with no individual researcher responsible for performing  
17 either the 4AT or the CAM; that is, each researcher performed approximately equal numbers  
18 of the 4AT and the CAM. The two assessments took place strictly within a maximum of two  
19 hours of each other, with a target interval of 15 minutes. Researchers were blinded to each  
20 other's assessments, that is, reference standard results were not available to performers of  
21 the index and comparator tests, and vice versa. The design of either 4AT or CAM rather  
22 than both 4AT and CAM being performed by each participant was chosen to avoid burden  
23 on participants, and because the CAM testing process is longer than the 4AT and

information from the CAM process could influence scoring of the 4AT; some influence of 4AT item scores on the CAM could also be possible.

### **Statistical Analysis**

All analyses were performed using SAS version 9.3 (SAS Institute Inc., Cary, NC, United States).

### **Primary objective**

We calculated positive and negative predictive values, sensitivity and specificity for 4AT versus the reference standard. We reported the area under the receiver operating characteristic (ROC) curve and its 95% confidence interval (CI) for the 4AT.

### **Secondary objective**

Comparison of 4AT and CAM: we calculated positive predictive values (PPV) and negative predictive values (NPV), sensitivity and specificity (with exact binomial 95% CI) for CAM and 4AT, and estimated the difference (4AT minus CAM) for each, assessing statistical significance of differences using Fisher's exact test. The area under the ROC curve could not be calculated for the CAM as the outcome is binary. The overall performance of 4AT and CAM were each summarised using Youden's Index (sensitivity minus false positive rate) and the diagnostic odds ratio of sensitivity to specificity.

### **Subgroup analyses**

Predefined subgroup analyses assessed the impact of (a) time from presentation to recruitment (analysing those tested before or after 4h (ED) or 24h (medical admissions)) for

4AT, and (b) time between index test and reference standard (analysing those tested within 30 minutes compared to those tested later) for both 4AT and CAM.

### **Sensitivity analyses**

We performed predefined sensitivity analyses where the reference standard was indeterminate by defining delirium as present, and then absent. We also performed a post hoc sensitivity analysis by using the initial delirium classification recorded by the researcher at the time of the original bedside assessment (which was performed to inform clinical staff at that time). A further post-hoc sensitivity analysis assumed that any patient with a missing result for the index test (4AT or CAM) had delirium.

### **Missing data**

If data were missing for the reference standard assessment, CAM or 4AT, or if the reference standard assessment did not yield a clear diagnosis, data from these individuals were removed from statistical analysis.

### **Sample size**

We planned to randomise 900 patients, 450 to assessment by 4AT and 450 to CAM. For each of 4AT and CAM the width of the two-sided 95% confidence interval for specificity would be up to  $\pm 0.050$ ; and for sensitivity, up to  $\pm 0.120$ . The secondary objective comparing 4AT and CAM would have 83% power to detect a difference in specificity of 0.10 and 80%



power to detect a difference in sensitivity of 0.22, for a 5% two-sided significance level and analysis by continuity corrected chi-squared test.

## Results

Study recruitment commenced on 19 October 2015, and was completed on 30 December 2016, with final follow-up data collection and locking of the database on 29 June 2017. 4,928 patients were eligible, from whom 843 individuals (17.1%) were recruited across the three sites, and two withdrew before data collection, leaving 841 with data for analysis of whom 19 withdrew, 3 lost contact, 32 were classified as indeterminate from the reference standard data, and 2 had a missing outcome. Therefore, 785 individuals were included in the analyses (Figure 3). Recruitment did not reach the target of 900 through a combination of a lower than expected rate of recruitment and a limit to the available recruitment period. However, the number recruited allowed for adequate power to test the main hypotheses as confirmed by the study statisticians and the Trial Steering Committee.

**Figure 3. STARD diagram of flow of participants through the study (total across all three sites)**

Reference standard delirium prevalence was 12.1% (n=95 of 785). Individuals with delirium were older and were more likely to have dementia as documented through the clinical records or through the informant questionnaire (Table 1). Baseline characteristics for those randomised to the 4AT or CAM are shown in Additional Table 1. Reference standard delirium prevalence in those who had a valid 4AT assessment was 12.5% (n=49 of 392), and in those who had a valid CAM assessment was 10.9% (n=42 of 384). Delirium prevalence using the 4AT only as a diagnostic test was 14.3% (n=56 of 392) and for CAM only as a diagnostic test was 4.7% (n=18 of 384).

**Table 1. Baseline demographic and clinical characteristics stratified by reference standard delirium status**

	Total (N=785)	Delirium Present (N=95)	Delirium Absent (N=690)	P value
<b>Age (Years)</b>				
Mean (SD)	81.4 (6.4)	83.5 (6.9)	81.1 (6.3)	0.0007
Median [Q1-Q3]	81.0 [77.0-86.0]	84.0 [78.0-89.0]	81.0 [77.0-86.0]	
<b>Gender</b>				
Male, n (%)	349 (44.5%)	34 (35.8%)	315 (45.7%)	0.0697
Female, n (%)	436 (55.5%)	61 (64.2%)	375 (54.3%)	
<b>Dementia Diagnosis and/or IQCODE<sup>a</sup> ≥ 3.44<sup>a</sup></b>				
Yes, n (%)	111 (14.2%)	43 (45.3%)	68 (9.9%)	<0.0001
No, n (%)	673 (85.5%)	52 (54.7%)	621 (90.1%)	
Missing, * n (%)	1 (0.1%)	0 (0.0%)	1 (0.1%)	
<b>Location of First Assessment</b>				
Emergency Department, n (%)	53 (6.8%)	10 (10.5%)	43 (6.2%)	0.2624
Acute General Medical Ward, n (%)	665 (84.7%)	76 (80.0%)	589 (85.4%)	
Hospital Ward, n (%)	67 (8.5%)	9 (9.5%)	58 (8.4%)	

P-value from chi-squared (categorical variables) or t-test (continuous).

\*Missing category not included in chi-squared test.

<sup>a</sup>IQCODE is Informant Questionnaire for Cognitive Impairment in the Elderly.

## Diagnostic test accuracy of 4AT and CAM

The main diagnostic test accuracy results for the 4AT and CAM are shown in Table 2. At a 4AT cut-off score for delirium of  $>3$ , the sensitivity was 76% (95% CI 61 to 87%) and the specificity was 94% (95% CI 92 to 97%). The performance at different cut-off scores is shown in Additional Table 2. The area under the ROC curve for the 4AT was 0.90 (95% CI 0.84 to 0.96) (Figure 4). The CAM had a sensitivity of 40% (95% CI 26 to 57%) and a specificity of 100% (95% CI 98 to 100%).

## Figure 4. Receiver Operator Characteristic Curve for 4AT diagnostic accuracy

**Table 2. Diagnostic test accuracy of the 4AT the CAM for diagnosis of delirium (defined by reference standard assessment)**

	<b>Sensitivity</b>	<b>Specificity</b>	<b>Positive Predictive Value</b>	<b>Negative Predictive Value</b>	<b>Youden's Index</b>
4AT (>3), <i>n</i> (95% CI)	76% (61 to 87%)	94% (92 to 97%)	66% (52 to 78%)	96% (94 to 98%)	0.70
CAM Positive , <i>n</i> (95% CI)	40% (26 to 57%)	100% (98 to 100%)	94% (73 to 100%)	93% (90 to 96%)	0.40
Difference in Proportions	36% (15 to 53%)	-6% (-14 to 2%)	-28% (-53 to -2%)	3% (-4 to 11%)	
<i>P</i> value	0.0012	<0.0001	0.0297	0.0629	

Numbers are estimates (95% CI). Youden's Index is equal to sensitivity+specificity-1, a value of zero indicates no value, and a value of 1 indicates a perfect test. The Difference in Proportions is 4AT-CAM for each of the tabulated measures of diagnostic accuracy, accompanied by the corresponding P-value from the Fisher's exact test comparing proportions. Abbreviations: CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value.

## Subgroup analyses

There was no statistically significant difference in the diagnostic test accuracy of the 4AT between those recruited early and those recruited later after initial presentation (Fisher's Exact Test p-values: sensitivity p=0.19, specificity p=0.75, PPV p=0.47, NPV p=0.24).

There was no statistically significant difference in performance of either test regardless of whether or not it was performed within 30 minutes of the reference standard (Fisher's Exact Test p-values: sensitivity p=0.16, specificity p=0.24, PPV p=1.00, NPV p=0.56).

## Sensitivity analyses

### Indeterminate reference standard

Assuming delirium was present for all indeterminate reference standards (N=32) reduced the sensitivity of both the 4AT and CAM: 64% (95% CI 52 to 76%), and 33% (95% CI 21 to 47%), respectively (Additional Table 3). Assuming delirium was absent for all indeterminate reference standards did not substantially alter the diagnostic accuracy of the 4AT or CAM (Additional Table 4).

### Delirium reference standard

Using the researchers' initial reference standard assessment of delirium, the sensitivity of the 4AT was 83% (95% CI 70 to 93%) and the specificity was 94% (95% CI 91 to 96%). The sensitivity (40%; 95% CI 25 to 56%) and specificity (99%; 95% CI 98 to 100%) of the CAM did not change substantially.

## Missing index test

If delirium was scored as present where the index test result was missing, this did not substantially alter the diagnostic test accuracy of the 4AT or CAM (Additional Table 5).

## Discussion

This study found that the 4AT had a sensitivity of 76% and a specificity of 94% for delirium as assessed independently by a reference standard. The area under the ROC curve was high at 0.90. These findings, in a relatively large, STARD-compliant study, provide support additional to the existing literature for the use of the 4AT as a delirium assessment instrument in clinical practice which has acceptable overall diagnostic test accuracy. The study also found that the CAM showed lower sensitivity than the 4AT, at 40%, with higher specificity at 100%. This is the first randomised comparison of two of the most widely used delirium assessment tools in clinical practice, and thus is informative for researchers with respect to their respective performance under the same study conditions.

The diagnostic test accuracy of the 4AT was broadly similar to the existing studies [25, 33-39], albeit with lower sensitivity and higher specificity than most prior studies. The difference in the sensitivity results may reflect differences in study population, the reference standard assessment, and recruitment processes. One prior study found higher sensitivity (87%) and lower specificity (70%) and a similar area under the ROC curve of 0.84 in an unselected consecutive clinical sample using a design that did not require consenting

(N=434)[35]. The 4AT involves a degree of subjectivity with respect to the assessment of level of alertness; raters are asked to rate this in a binary fashion, that is, as abnormal or normal. The reference standard assessment involves a more detailed approach to assessment of arousal involving the three different tools: the Observational Scale of Level of Arousal, the Richmond Agitation-Sedation Scale, and the arousal element of the DelApp smartphone test of attention and arousal. It is possible that the simpler binary assessment tended to a lower level of positive score than the more complex and nuanced reference standard assessment process. Additionally, the bedside element of the 4AT (items 1-3) usually takes less than one minute, as compared to around 20 minutes for the reference standard assessment. This give more opportunity in the reference standard assessment for the observation of reduced arousal as well as fluctuation of symptoms. Further planned analyses of the present dataset will explore the relationships of individual test components of the 4AT (and the CAM) to the overall test score and components of the reference standard.

In this study the CAM showed very high specificity and modest sensitivity for delirium. The high specificity is aligned with prior studies, the vast majority of which have found specificities of over 90%. The sensitivity of 40% was lower than in the majority of published studies. However, unlike with specificity, the literature shows notable heterogeneity in findings with respect to CAM sensitivity, with several studies also showing lower sensitivities for the CAM [48, 50, 51, 57, 65, 67]. Differences in study populations, eligibility criteria (e.g. exclusion of drowsy patients unable to produce speech), the interview and cognitive testing performed, the training provided (this is variably described in the literature), and the background and experience of the raters may all play a role in the variability of findings [48, 59, 81]. The CAM involves binary, subjective bedside judgements of inattention, disorganised thinking, and level of consciousness; such



1 judgements are more open to variability between raters compared to objective scoring [72,  
2 75]. Another possible source of reduced sensitivity in some studies is that the CAM  
3 algorithm generates a negative score if disorganised thinking is not ascertained (that is, if  
4 ‘rambling, irrelevant or incoherent speech’ [40] is not judged to be present) and if the level  
5 of consciousness is judged to be normal, though the patient may have inattention and other  
6 cognitive deficits and thus meet DSM-IV or DSM-5 criteria for delirium. Similarly, if  
7 inattention is not judged to be present but there is altered level of consciousness the CAM  
8 algorithm will generate a negative score.

9 This study had several strengths. Each participant was randomised to perform either the 4AT  
10 or CAM under the same study conditions, with the reference standard being performed  
11 independently by a different researcher. This is of interest given that the 4AT and the CAM  
12 are two of the most commonly-used tools internationally. Researchers were formally trained  
13 in use of the CAM and the reference standard assessment. The reference assessment  
14 involved gathering information from the DRS–R98, several tests of cognition and also level  
15 of arousal. Neufeld and colleagues [82] found substantial variability in delirium reference  
16 standard assessments used in diagnostic accuracy studies of delirium assessment tools, with  
17 many not using cognitive testing as part of the assessment process. The present study had  
18 limited exclusion criteria, allowing patients with a wide spectrum of level of severity to be  
19 approached, including patients with severely reduced level of arousal. This is pertinent  
20 because reduced level of arousal is common in emergency admissions; in one study of  
21 clinically-collected data from 35,585 consecutive, unselected acute medical admissions  
22 aged >15, 7.6% of patients had reduced level of arousal above the level of coma, and in  
23 older populations the prevalence is are higher [73, 83-85]. Given the close relationship of  
24 reduced arousal with delirium [29, 32, 73, 85, 86] it is important that studies of delirium

assessment instruments include the full spectrum of patients with reduced arousal (excluding coma). The study was relatively large, and multicentre. The protocol was published in advance of database lock and analysis, and the study reporting adhered to the STARD guidelines.

Some limitations of this study should be acknowledged. In this study only 17% of those eligible for recruitment were recruited, mostly due to patients declining to participate or no person available to provide proxy consent. The delirium rate was 12.1% according to the reference standard; prior studies have estimated that the prevalence of delirium in patients aged 70 or above at the early stages of hospital admission likely ranges from 10 to 20% [87]. The recruitment process, which required consenting (often from a proxy), may have led to a sample with a moderately lower delirium prevalence than in clinical populations. This is a known limitation of delirium studies requiring consent [88]. Most patients with delirium lack capacity, and in the context of the present study this necessitated proxy consent and an informant to score the acute change items in the 4AT and CAM. In clinical practice, the acute change item might be informed by staff knowledge of the patient, or not scored if no such information exists (though an overall positive score is still possible on the 4AT because of the scoring procedure for items (A), (B), and (C); this differs from the process that was required in the study. With respect to the reference standard, it is possible that objective assessments recorded and interpreted for this did not fully capture the researcher's interaction with the patient and thus the researcher's ascertainment of DSM-IV delirium features. Results from the sensitivity analysis using bedside reference standard diagnosis support this possibility, showing higher rates of both sensitivity (83%) and specificity (94%) if the researcher's initial assessment was used. We aimed to ascertain dementia status but it is possible that some patients had dementia but this was undiagnosed and the IQCODE was

1 unavailable. The number of patients with known dementia was too low to allow analysis of  
2 performance of the 4AT or CAM in patients with and without dementia. Finally, it is  
3 possible that researcher bias may have influenced the conduct or scoring of the different  
4 index assessments (4AT or CAM) because the 4AT was designed in one of the sites of the  
5 study and involved AMJM. However none of the researchers collecting data was involved  
6 in development of the 4AT, the CAM was performed by researchers trained in its use as  
7 advised in the CAM instruction manual, and the reference standard was administered by  
8 researchers blind to the identity or results of the index tests.

9 Future studies could seek to compare performance of the 4AT with other short delirium  
10 assessment tests, such as the Single Question in Delirium (SQiD)[61], the Delirium Triage  
11 Screen [89], the brief CAM (bCAM)[89, 90], the 3D-CAM [91] and the Simple Query for  
12 Easy Evaluation of Consciousness (SQeeC)[65]. Studies could also evaluate the value of  
13 the individual items of the 4AT. This is an important issue because though ideally informant  
14 history is used to make a diagnosis of delirium, in a substantial proportion of patients such  
15 history is not available at the point of initial assessment or even during the inpatient stay  
16 [35, 38]. Additionally, the extent of real-world use in large clinical datasets including rates  
17 of positive scores should be evaluated. For example the 4AT is mandated to assess for post-  
18 operative delirium in all acute hip fracture patients in the National Health Service in  
19 England, Wales, and Northern Ireland; in 2017 86% of 63,471 patients were assessed with  
20 the 4AT, with 25% showing a positive score [92]. This is possibly an underestimate of post-  
21 operative delirium rates, but suggests that the 4AT is embedded in routine practice and likely  
22 detecting the majority of delirium. Further reporting of use of the 4AT and other tools in  
23 other large clinical datasets will be informative in determining feasibility outside of research  
24 studies.

## Conclusions

The 4AT showed moderate sensitivity, high specificity, and good overall diagnostic performance. In the present study the 4AT showed higher sensitivity than the CAM, and slightly lower specificity under the same study conditions. The CAM has been evaluated by multiple validation studies and while many of these studies show high sensitivity, many also show that sensitivity tends to be lower where raters are not fully trained in the CAM, or who lack specialist training in psychiatric assessment. Taken as a whole, the 4AT validation studies suggest that it has comparable performance to the CAM when the CAM is being performed by trained raters. In terms of its brevity, lack of need for training, and comparable performance the 4AT can therefore reasonably be used as an assessment tool for delirium, particularly in clinical settings in which there is limited time, and in which staff involved in delirium detection cannot undergo the substantial special training required for use of the CAM. Additionally, the 4AT can be scored if no informant history is available at the time of assessment, and if arousal is impaired such that cognitive testing is not possible, which occurs in a substantial proportion of delirium assessments [35, 38, 65]. Given that acutely altered arousal is a highly specific indicator of delirium [29, 30, 72, 73, 85, 93-95] and that it often indicates a poor prognosis [83], a rapid provisional diagnosis of delirium with appropriate action in the absence of an external informant history is reasonable. The 4AT is designed to be able to yield a positive score in patients too unwell to undergo interview or cognitive testing [31, 32]; this facilitates implementation and higher completion rates in clinical practice. It is important to note, however, that as with all short detection tools, formal diagnosis of delirium in clinical practice requires assessment by a suitably-qualified member of staff.

## Figure legends

**Figure 1 legend: [no legend]**

**Figure 2 legend:** ED = Emergency Department; MOE = Medicine of the Elderly; 4AT = 4As Tests; CAM = Confusion Assessment Method

**Figure 3 legend: [no legend]**

**Figure 4 legend:** 4AT scores range from 0-12. The cut-point of >3 is used in the scoring scheme to denote likely delirium. The 4AT scores are considered against the reference standard delirium assessment.

**Additional Table 1: Baseline Characteristics by Index Test (4AT or CAM).** Legend: Numbers are n (%) or mean (SD).

**Additional Table 2: Performance of various cut points of 4AT for diagnosis of delirium.** Legend: Numbers are estimates (95% CI). Abbreviations: CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value. Youden's Index is equal to sensitivity+specificity-1, a value of zero indicates no value, and a value of 1 indicates a perfect test.

**Additional Table 3: Sensitivity analysis of diagnostic test accuracy of 4AT versus CAM for diagnosis of delirium assuming all indeterminates are delirium present.** Legend: Numbers are estimate (95% CI). Difference in proportions is for 4AT-CAM. Abbreviations: CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value; OR, odds ratio. Youden's Index is equal to sensitivity+specificity-1, a value of zero indicates no value, and a value of 1 indicates a perfect test.

**Additional Table 4: Sensitivity analysis of diagnostic test accuracy of 4AT versus CAM for diagnosis of delirium assuming all indeterminates are delirium absent.** Legend:

Numbers are estimate (95% CI). Difference in proportions is for 4AT-CAM. Abbreviations: CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value; OR, odds ratio. Youden's Index is equal to sensitivity+specificity-1, a value of zero indicates no value, and a value of 1 indicates a perfect test.

**Additional Table 5: Diagnostic test accuracy of 4AT versus CAM for diagnosis of delirium, assuming test scored delirium present for those with a missing 4AT or CAM score.** Legend: Numbers are estimate (95% CI). Difference in proportions is for 4AT-CAM.

Abbreviations: CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value; OR, odds ratio. Youden's Index is equal to sensitivity+specificity-1, a value of zero indicates no value, and a value of 1 indicates a perfect test.

## List of abbreviations

3D-CAM: 3-Minute Diagnostic Assessment for Delirium using the Confusion Assessment Method

4AT: 4 'A's Test

bCAM: Brief Confusion Assessment Method

CAM: Confusion Assessment Method

CI: confidence interval

- 1 DSM-III-R: Diagnostic and Statistical Manual, 3<sup>rd</sup> edition, revised
- 2
- 3
- 4 2 DSM-IV: Diagnostic and Statistical Manual, 4<sup>th</sup> edition
- 5
- 6
- 7 3 DSM-5: Diagnostic and Statistical Manual, 5<sup>th</sup> edition
- 8
- 9
- 10
- 11 4 ED: Emergency Department
- 12
- 13
- 14 5 IQCODE: Informant Questionnaire on Cognitive Decline in the Elderly
- 15
- 16
- 17
- 18 6 ISRCTN: International standard randomised controlled trial number
- 19
- 20
- 21
- 22 7 NPV: negative predictive value
- 23
- 24
- 25 8 NIHR HTA: National Institute of Health Research Health Technology Assessment
- 26
- 27
- 28 9 Programme
- 29
- 30
- 31
- 32 10 NHS: National Health Service
- 33
- 34
- 35 11 PPV: positive predictive value
- 36
- 37
- 38
- 39 12 REC: research ethics committee
- 40
- 41
- 42
- 43 13 ROC: receiver operating characteristic
- 44
- 45
- 46 14 SQiD: Single Question in Delirium
- 47
- 48
- 49
- 50 15 SqueC: Simple Query for Easy Evaluation of Consciousness
- 51
- 52
- 53
- 54 16 STARD: Standards for Reporting Diagnostic Accuracy
- 55
- 56
- 57
- 58
- 59

## Declarations

### Ethics approval and consent to participate

This study was granted ethical approval prior to data collection in Scotland (Scotland A NHS Research Ethics Committee REC 15/SS/0071) and England (Yorkshire and The Humber – Bradford Leeds NHS Research Ethics Committee REC 15/YH/0317).

### Consent for publication

Not applicable

### Availability of data and materials

Analyses of the data in this study are still ongoing. We shall make fully anonymised data available on the website <https://datashare.is.ed.ac.uk/> in an estimated one year from the publication of this manuscript.

### Competing interests

AMJM led the design of the 4AT in 2011 (with others, see [www.the4AT.com](http://www.the4AT.com)); note that 4AT is free to download and use. SDS and AA provided comments on its development. SG is chair of the NIHR HTA Clinical Evaluation and Trials Board and member of the NIHR HTA Funding Board Policy Group. The other authors declare no competing interests.



## **Funding**

Funder: National Institute of Health Research, ref. 11/143/01. The research proposal was developed by the authors in response to a commissioning call from the funder for a study to compare a new delirium assessment tool to the Confusion Assessment Method (CAM). The funder had no other role in the study design or conduct of the study.

## **Authors' contributions**

All authors helped draft the paper and critiqued the paper for important intellectual content. SDS provided expertise in geriatric medicine, and made a substantial contribution to protocol design and statistical analysis, and training and supervision of staff. CF provided expertise in dementia, and made a substantial contribution to protocol design. MG made a substantial contribution to the design and analysis of the study. NS provided expertise in psychiatry and made a substantial contribution to protocol design. SG provided expertise in emergency medicine, and made a substantial contribution to protocol design, staff training and supervision. JY provided expertise in geriatric medicine and made a substantial contribution to protocol design, staff training and supervision. AA provided expertise in geriatric medicine and made a substantial contribution to protocol design. JH made a substantial contribution to the design and analysis of the study. AG provided expertise in emergency medicine, and made a substantial contribution to the design of the study, staff training and supervision. AM, Jills, and PB provided expertise in nursing, made a substantial contribution to study design and participated in patient recruitment and testing, and in staff training. ZT provided expertise in psychology, and made a substantial contribution to design of the reference standard assessment, and to statistical analysis. JB provided expertise in study management, and made a substantial contribution to protocol

1 design and staff training and supervision. JacqS was a trial statistician and made a substantial  
2 contribution to the development of the statistical analysis plan and undertook the statistical  
3 analyses. CW was a trial statistician, and co-designed the protocol, led the design of the  
4 statistical analysis plan, and oversaw the statistical analyses. AMJM conceived and planned  
5 the study and acted as study lead, making substantial contributions to design of the reference  
6 standard assessment, statistical analysis and training of research staff.

## 7 **Acknowledgements**

8 The authors would like to thank all the patients and carers who have taken the time to take  
9 part in the study. We would also like to thank all the NHS staff who supported the study in  
10 the three sites. The authors acknowledge the help of the Trial Steering Committee. We thank  
11 all the administrative staff in the three study sites for their support. We thank Dr Valentina  
12 Assi for contributing to the statistical analyses. We acknowledge the help of the Edinburgh  
13 Clinical Trials Unit. We thank Miranda Odam, Mia Paderanga, and Louise Ross from the  
14 Emergency Medicine Research Group of Edinburgh (EMERGE). We thank the research  
15 nurses and other research staff in the Clinical Research Facilities who took part in the  
16 recruitment and testing.

# References

1. Reynish EL, Hapca SM, De Souza N, Cvorov V, Donnan PT, Guthrie B: **Epidemiology and outcomes of people with dementia, delirium, and unspecified cognitive impairment in the general hospital: prospective cohort study of 10,014 admissions.** *BMC Medicine* 2017, **15**(1):140.
2. Pendlebury ST, Lovett NG, Smith SC, Dutta N, Bendon C, Lloyd-Lavery A, Mehta Z, Rothwell PM: **Observational, longitudinal study of delirium in consecutive unselected acute medical admissions: age-specific rates and associated factors, mortality and re-admission.** *BMJ open* 2015, **5**(11):e007808.
3. Marcantonio ER: **Delirium in Hospitalized Older Adults.** *N Engl J Med* 2017, **377**(15):1456-1466.
4. Oh ES, Fong TG, Hsieh TT, Inouye SK: **Delirium in Older Persons: Advances in Diagnosis and Treatment.** *JAMA* 2017, **318**(12):1161-1174.
5. Goldberg SE, Whittamore KH, Harwood RH, Bradshaw LE, Gladman JR, Jones RG: **The prevalence of mental health problems among older adults admitted as an emergency to a general hospital.** *Age Ageing* 2012, **41**(1):80-86.
6. The Royal College of Psychiatrists: **Who Cares Wins.** London: Royal College of Psychiatrists; 2005.
7. American Psychiatric Association: **Diagnostic and statistical manual of mental disorders : DSM-IV-TR**, 4th edn. Washington, DC; 2000.
8. American Psychiatric Association: **Diagnostic and Statistical Manual of Mental Disorders**, 5th edn. Washington, DC; 2013.
9. Davis DH, Muniz-Terrera G, Keage HA, Stephan BC, Fleming J, Ince PG, Matthews FE, Cunningham C, Ely EW, MacLulich AM *et al*: **Association of Delirium With Cognitive Decline in Late Life: A Neuropathologic Study of 3 Population-Based Cohort Studies.** *JAMA Psychiatry* 2017, **74**(3):244-251.
10. MacLulich AMJ, Beaglehole A, Hall RJ, Meagher DJ: **Delirium and long-term cognitive impairment.** *International Review of Psychiatry* 2009, **21**(1):30-42.
11. Witlox J, Eurelings LSM, de Jonghe JFM, Kalisvaart KJ, Eikelenboom P, van Gool WA: **Delirium in Elderly Patients and the Risk of Postdischarge Mortality, Institutionalization, and Dementia: A Meta-analysis.** *JAMA* 2010, **304**(4):443-451.

12. Han JH, Shintani A, Eden S, Morandi A, Solberg LM, Schnelle J, Dittus RS, Storrow AB, Ely EW: **Delirium in the emergency department: an independent predictor of death within 6 months.** *Ann Emerg Med* 2010, **56**(3):244-252 e241.
13. Burton JK, Guthrie B, Hapca SM, Cvorov V, Donnan PT, Reynish EL: **Living at home after emergency hospital admission: prospective cohort study in older adults with and without cognitive spectrum disorder.** *BMC Medicine* 2018, **16**(1):231.
14. Partridge JS, Martin FC, Harari D, Dhesi JK: **The delirium experience: what is the effect on patients, relatives and staff and what can be done to modify this?** *Int J Geriatr Psychiatry* 2013, **28**(8):804-812.
15. Martins S, Pinho E, Correia R, Moreira E, Lopes L, Paiva JA, Azevedo L, Fernandes L: **What effect does delirium have on family and nurses of older adult patients?** *Aging Ment Health* 2018, **22**(7):903-911.
16. Racine AM, D'Aquila M, Schmitt EM, Gallagher J, Marcantonio ER, Jones RN, Inouye SK, Schulman-Green D, Group BS: **Delirium Burden in Patients and Family Caregivers: Development and Testing of New Instruments.** *Gerontologist* 2019, **59**(2):327-337.
17. Collins N, Blanchard MR, Tookman A, Sampson EL: **Detection of delirium in the acute hospital.** *Age Ageing* 2010, **39**(1):131-135.
18. Han JH, Zimmerman EE, Cutler N, Schnelle J, Morandi A, Dittus RS, Storrow AB, Ely EW: **Delirium in older emergency department patients: recognition, risk factors, and psychomotor subtypes.** *Academic Emergency Medicine* 2009, **16**(3):193-200.
19. Traynor V, Cordato N, Burns P, Xu Y, Britten N, Duncan K, DeVries L, McKinnon C: **Is delirium being detected in emergency?** *Australas J Ageing* 2016, **35**(1):54-57.
20. Bellelli G, Nobili A, Annoni G, Morandi A, Djade CD, Meagher DJ, MacLulich AM, Davis D, Mazzone A, Tettamanti M *et al*: **Under-detection of delirium and impact of neurocognitive deficits on in-hospital mortality among acute geriatric and medical wards.** *Eur J Intern Med* 2015, **26**(9):696-704.
21. Stelmokas J, Gabel N, Flaherty JM, Rayson K, Tran K, Anderson JR, Bieliauskas LA: **Delirium Detection and Impact of Comorbid Health Conditions in a Post-Acute Rehabilitation Hospital Setting.** *PLoS One* 2016, **11**(11):e0166754.
22. Davis D, MacLulich A: **Understanding barriers to delirium care: a multicentre survey of knowledge and attitudes amongst UK junior doctors.** *Age Ageing* 2009, **38**(5):559-563.
23. Fisher JM, Gordon AL, MacLulich AM, Tullo E, Davis DH, Blundell A, Field RH, Teodorczuk A: **Towards an understanding of why undergraduate teaching about delirium does not guarantee gold-standard practice--results from a UK national survey.** *Age Ageing* 2015, **44**(1):166-170.

24. Teodorczuk A, Reynish E, Milisen K: **Improving recognition of delirium in clinical practice: a call for action.** *BMC Geriatr* 2012, **12**:55.
25. Bellelli G, Morandi A, Davis DH, Mazzola P, Turco R, Gentile S, Ryan T, Cash H, Guerini F, Torpilliesi T *et al*: **Validation of the 4AT, a new instrument for rapid delirium screening: a study in 234 hospitalised older people.** *Age Ageing* 2014, **43**(4):496-502.
26. **The 4 "A"s Test** [[www.the4AT.com](http://www.the4AT.com)]. Accessed 1 September 2014.
27. Shenkin SD, Fox C, Godfrey M, Siddiqi N, Goodacre S, Young J, Anand A, Gray A, Smith J, Ryan T *et al*: **Protocol for validation of the 4AT, a rapid screening tool for delirium: a multicentre prospective diagnostic test accuracy study.** *BMJ Open* 2018, **8**(2):e015572.
28. Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI: **Clarifying confusion: the confusion assessment method. A new method for detection of delirium.** *Annals of Internal Medicine* 1990, **113**(12):941-948.
29. Tieges Z, McGrath A, Hall RJ, MacLulich AM: **Abnormal level of arousal as a predictor of delirium and inattention: an exploratory study.** *American Journal of Geriatric Psychiatry* 2013, **21**(12):1244-1253.
30. Chester JG, Harrington MB, Rudolph JL, Grp VADW: **Serial administration of a modified richmond agitation and sedation scale for delirium screening.** *Journal of Hospital Medicine* 2012, **7**(5):450-453.
31. Yates C, Stanley N, Cerejeira JM, Jay R, Mukaetova-Ladinska EB: **Screening instruments for delirium in older people with an acute medical illness.** *Age Ageing* 2009, **38**(2):235-237.
32. European Delirium Association, American Delirium Society: **The DSM-5 criteria, level of arousal and delirium diagnosis: inclusiveness is safer.** *BMC Medicine* 2014, **12**:141.
33. Lees R, Corbet S, Johnston C, Moffitt E, Shaw G, Quinn TJ: **Test accuracy of short screening tests for diagnosis of delirium or cognitive impairment in an acute stroke unit setting.** *Stroke* 2013, **44**(11):3078-3083.
34. Kuladee S, Prachason T: **Development and validation of the Thai version of the 4 'A's Test for delirium screening in hospitalized elderly patients with acute medical illnesses.** *Neuropsychiatr Dis Treat* 2016, **12**:437-443.
35. Hendry K, Quinn TJ, Evans J, Scortichini V, Miller H, Burns J, Cunningham A, Stott DJ: **Evaluation of delirium screening tools in geriatric medical inpatients: a diagnostic test accuracy study.** *Age Ageing* 2016, **45**(6):832-837.

36. De J, Wand AP, Smerdely PI, Hunt GE: **Validating the 4A's test in screening for delirium in a culturally diverse geriatric inpatient population.** *Int J Geriatr Psychiatry* 2017, **32**(12):1322-1239.
37. Infante MT, Pardini M, Balestrino M, Finocchi C, Malfatto L, Bellelli G, Mancardi GL, Gandolfo C, Serrati C: **Delirium in the acute phase after stroke: comparison between methods of detection.** *Neurol Sci* 2017, **38**(6):1101-1104.
38. O'Sullivan D, Brady N, Manning E, O'Shea E, O'Grady S, N OR, Timmons S: **Validation of the 6-Item Cognitive Impairment Test and the 4AT test for combined delirium and dementia screening in older Emergency Department attendees.** *Age Ageing* 2018, **47**(1):61-68.
39. Saller T, MacLulich A, Schaher T, Crispin A, Neitzert R, Schule C, Von Dossow V, Hofmann-Kiefer KF: **Validation study of the 4 'A's Test (4AT) for delirium detection in post-anaesthesia care.** *Anaesthesia* 2019, Apr 30. [Epub]
40. Inouye SK: **The Short Confusion Assessment Method (Short CAM): Training Manual and Coding Guide.** Boston: Hospital Elder Life Program; 2014.
41. **Delirium: prevention, diagnosis and management** [<https://www.nice.org.uk/guidance/cg103>] Accessed 1 September 2014.
42. Wei LA, Fearing MA, Sternberg EJ, Inouye SK: **The Confusion Assessment Method: a systematic review of current usage.** *Journal of the American Geriatrics Society* 2008, **56**(5):823-830.
43. Wong CL, Holroyd-Leduc J, Simel DL, Straus SE: **Does this patient have delirium?: value of bedside instruments.** *JAMA* 2010, **304**(7):779-786.
44. Shi Q, Warren L, Saposnik G, Macdermid JC: **Confusion assessment method: a systematic review and meta-analysis of diagnostic accuracy.** *Neuropsychiatr Dis Treat* 2013, **9**:1359-1370.
45. De J, Wand AP: **Delirium Screening: A Systematic Review of Delirium Screening Tools in Hospitalized Patients.** *The Gerontologist* 2015, **55**(6):1079-1099.
46. van Velthuisen EL, Zwakhalen SM, Warnier RM, Mulder WJ, Verhey FR, Kempen GI: **Psychometric properties and feasibility of instruments for the detection of delirium in older hospitalized patients: a systematic review.** *Int J Geriatr Psychiatry* 2016, **31**(9):974-989.
47. Inouye SK, Vandyck CH, Alessi CA, Balkin S, Siegal AP, Horwitz RI: **Clarifying Confusion - the Confusion Assessment Method - A New Method for Detection of Delirium.** *Annals of Internal Medicine* 1990, **113**:941-948.
48. Rockwood K, Cosway S, Stolee P, Kydd D, Carver D, Jarrett P, O'Brien B: **Increasing the recognition of delirium in elderly patients.** *J Am Geriatr Soc* 1994, **42**(3):252-256.

49. Farrell KR, Ganzini L: **Misdiagnosing delirium as depression in medically ill elderly patients.** *Arch Intern Med* 1995, **155**(22):2459-2464.
50. Pompei P, Foreman M, Cassel CK, Alessi C, Cox D: **Detecting delirium among hospitalized older patients.** *Arch Intern Med* 1995, **155**(3):301-307.
51. Rolfson DB, McElhaney JE, Jhangri GS, Rockwood K: **Validity of the confusion assessment method in detecting postoperative delirium in the elderly.** *International Psychogeriatrics* 1999, **11**(4):431-438.
52. Fabbri RM, Moreira MA, Garrido R, Almeida OP: **Validity and reliability of the Portuguese version of the Confusion Assessment Method (CAM) for the detection of delirium in the elderly.** *Arq Neuropsiquiatr* 2001, **59**(2-A):175-179.
53. Laurila JV, Pitkala KH, Strandberg TE, Tilvis RS: **Confusion assessment method in the diagnostics of delirium among aged hospital patients: would it serve better in screening than as a diagnostic instrument?** *Int J Geriatr Psychiatry* 2002, **17**(12):1112-1119.
54. Gonzalez M, de Pablo J, Fuente E, Valdes M, Peri JM, Nomdedeu M, Matrai S: **Instrument for detection of delirium in general hospitals: adaptation of the confusion assessment method.** *Psychosomatics* 2004, **45**(5):426-431.
55. Gaudreau JD, Gagnon P, Harel F, Tremblay A, Roy MA: **Fast, systematic, and continuous delirium assessment in hospitalized patients: the nursing delirium screening scale.** *J Pain Symptom Manage* 2005, **29**(4):368-375.
56. Leung J, Leung V, Leung CM, Pan PC: **Clinical utility and validation of two instruments (the Confusion Assessment Method Algorithm and the Chinese version of Nursing Delirium Screening Scale) to detect delirium in geriatric inpatients.** *Gen Hosp Psychiatry* 2008, **30**(2):171-176.
57. Radtke FM, Franck M, Schneider M, Luetz A, Seeling M, Heinz A, Wernecke KD, Spies CD: **Comparison of three scores to screen for delirium in the recovery room.** *Br J Anaesth* 2008, **101**(3):338-343.
58. Hestermann U, Backenstrass M, Gekle I, Hack M, Mundt C, Oster P, Thomas C: **Validation of a German version of the Confusion Assessment Method for delirium detection in a sample of acute geriatric patients with a high prevalence of dementia.** *Psychopathology* 2009, **42**(4):270-276.
59. Ryan K, Leonard M, Guerin S, Donnelly S, Conroy M, Meagher D: **Validation of the confusion assessment method in the palliative care setting.** *Palliat Med* 2009, **23**(1):40-45.
60. Radtke FM, Franck M, Schust S, Boehme L, Pascher A, Bail HJ, Seeling M, Luetz A, Wernecke KD, Heinz A *et al*: **A comparison of three scores to screen for delirium on the surgical ward.** *World J Surg* 2010, **34**(3):487-494.

- 1 61. Sands MB, Dantoc BP, Hartshorn A, Ryan CJ, Lujic S: **Single Question in Delirium (SQiD): testing its efficacy against psychiatrist interview, the Confusion Assessment Method and the Memorial Delirium Assessment Scale.** *Palliative Medicine* 2010, **24**(6):561-565.
- 2 62. Wongpakaran N, Wongpakaran T, Bookamana P, Pinyopornpanish M, Maneeton B, Lerttrakarnnon P, Uttawichai K, Jiraniramai S: **Diagnosing delirium in elderly Thai patients: utilization of the CAM algorithm.** *BMC Fam Pract* 2011, **12**:65.
- 3 63. Thomas C, Kreisel SH, Oster P, Driessen M, Arolt V, Inouye SK: **Diagnosing delirium in older hospitalized adults with dementia: adapting the confusion assessment method to international classification of diseases, tenth revision, diagnostic criteria.** *J Am Geriatr Soc* 2012, **60**(8):1471-1477.
- 4 64. Charoensak S, Thunmanurukkit A, Sittironnarit G, Sartra T: **Validity and reliability of the Thai version of the confusion assessment method.** *J Med Assoc Thai* 2014, **97**(1):113-117.
- 5 65. Lin HS, Eeles E, Pandey S, Pinsker D, Brasch C, Yerkovich S: **Screening in delirium: A pilot study of two screening tools, the Simple Query for Easy Evaluation of Consciousness and Simple Question in Delirium.** *Australas J Ageing* 2015, **34**(4):259-264.
- 6 66. Martins S, Lourenco C, Pinto-de-Sousa J, Conceicao F, Paiva JA, Simoes MR, Fernandes L: **Validation study of the European Portuguese version of the Confusion Assessment Method (CAM).** *International Psychogeriatrics* 2015, **27**(5):777-784.
- 7 67. Smulter N, Lingehall HC, Gustafson Y, Olofsson B, Engstrom KG: **Validation of the confusion assessment method in detecting postoperative delirium in cardiac surgery patients.** *Am J Crit Care* 2015, **24**(6):480-487.
- 8 68. Cohen JF, Korevaar DA, Altman DG, Bruns DE, Gatsonis CA, Hooft L, Irwig L, Levine D, Reitsma JB, de Vet HC *et al*: **STARD 2015 guidelines for reporting diagnostic accuracy studies: explanation and elaboration.** *BMJ Open* 2016, **6**(11):e012799.
- 9 69. Shenkin SD, Fox C, Godfrey M, Siddiqi N, Goodacre S, Young J, Anand A, Gray A, Smith J, Ryan T *et al*: **Protocol for validation of the 4AT, a rapid screening tool for delirium: a multicentre prospective diagnostic test accuracy study.** *BMJ Open* 2018, **8**(2):e015572.
- 10 70. Trzepacz PT, Mittal D, Torres R, Canary K, Norton J, Jimerson N: **Validation of the Delirium Rating Scale-revised-98: Comparison with the Delirium Rating Scale and the Cognitive Test for Delirium.** *Journal of Neuropsychiatry and Clinical Neurosciences* 2001, **13**(2):229-242.
- 11 71. Tieges Z, Brown LJ, MacLulich AM: **Objective assessment of attention in delirium: a narrative review.** *Int J Geriatr Psychiatry* 2014, **29**(12):1185-1197.



72. Tieges Z, Evans JJ, Neufeld KJ, MacLulich AM: **The neuropsychology of delirium: advancing the science of delirium assessment.** *Int J Geriatr Psychiatry* 2018, **33**(11):1501-1511.
73. Richardson SJ, Davis DHJ, Bellelli G, Hasemann W, Meagher D, Kreisel SH, MacLulich AMJ, Cerejeira J, Morandi A: **Detecting delirium superimposed on dementia: diagnostic accuracy of a simple combined arousal and attention testing procedure.** *International Psychogeriatrics* 2017, **29**(10):1585-1593.
74. Sessler CN, Gosnell MS, Grap MJ, Brophy GM, O'Neal PV, Keane KA, Tesoro EP, Elswick RK: **The Richmond Agitation-Sedation Scale: validity and reliability in adult intensive care unit patients.** *American Journal of Respiratory and Critical Care Medicine* 2002, **166**(10):1338-1344.
75. Simon SE, Bergmann MA, Jones RN, Murphy KM, Orav EJ, Marcantonio ER: **Reliability of a structured assessment for nonclinicians to detect delirium among new admissions to postacute care.** *Journal of the American Medical Directors Association* 2006, **7**(7):412-415.
76. Hart RP, Levenson JL, Sessler CN, Best AM, Schwartz SM, Rutherford LE: **Validation of a cognitive test for delirium in medical ICU patients.** *Psychosomatics* 1996, **37**(6):533-546.
77. Tieges Z, Stiobhairt A, Scott K, Suchorab K, Weir A, Parks S, Shenkin S, MacLulich A: **Development of a smartphone application for the objective detection of attentional deficits in delirium.** *International Psychogeriatrics* 2015, **27**(8):1251-1262.
78. Tang E, Lavery M, Weir A, Wilson ES, Walsh TS, Allerhand M, Parks S, MacLulich AMJ, Tieges Z: **Development and feasibility of a smartphone-based test for the objective detection and monitoring of attention impairments in delirium in the ICU.** *J Crit Care* 2018, **48**:104-111.
79. Rutter LM, Nouzova E, Stott DJ, Weir CJ, Assi V, Barnett JH, Clarke C, Duncan N, Evans J, Green S *et al*: **Diagnostic test accuracy of a novel smartphone application for the assessment of attention deficits in delirium in older hospitalised patients: a prospective cohort study protocol.** *BMC Geriatr* 2018, **18**(1):217.
80. Neufeld KJ, Nelliot A, Inouye SK, Ely EW, Bienvenu OJ, Lee HB, Needham DM: **Delirium diagnosis methodology used in research: a survey-based study.** *American Journal Of Geriatric Psychiatry* 2014, **22**(12):1513-1521.
81. Inouye SK, Foreman MD, Mion LC, Katz KH, Cooney LM: **Nurses' recognition of delirium and its symptoms - Comparison of nurse and researcher ratings.** *Archives of Internal Medicine* 2001, **161**:2467-2473.
82. Wozniak AW, Colantuoni EJ, Schreiber MP, Neufeld KJ, Needham DM: **Corticosteroids and transition to delirium in acute lung injury: multinomial**

- 1 **logistic regression analysis accounting for multiple States.** *Critical Care Medicine*  
2 2015, **43**(1):e23-24.
- 3 83. Todd A, Blackley S, Burton JK, Stott DJ, Ely EW, Tieges Z, MacLulich AMJ, Shenkin  
4 SD: **Reduced level of arousal and increased mortality in adult acute medical**  
5 **admissions: a systematic review and meta-analysis.** *BMC Geriatr* 2017, **17**(1):283.
- 6 84. Bellelli G, Mazzone A, Morandi A, Latronico N, Perego S, Zazzetta S, Mazzola P,  
7 Annoni G: **The Effect of an Impaired Arousal on Short- and Long-Term Mortality**  
8 **of Elderly Patients Admitted to an Acute Geriatric Unit.** *Journal of the American*  
9 *Medical Directors Association* 2016, **17**(3):214-219.
- 10 85. Morandi A, Han JH, Meagher D, Vasilevskis E, Cerejeira J, Hasemann W, MacLulich  
11 AM, Annoni G, Trabucchi M, Bellelli G: **Detecting Delirium Superimposed on**  
12 **Dementia: Evaluation of the Diagnostic Performance of the Richmond Agitation**  
13 **and Sedation Scale.** *Journal of the American Medical Directors Association* 2016,  
14 **17**(9):828-833.
- 15 86. Chester JG, Beth Harrington M, Rudolph JL: **Serial administration of a modified**  
16 **Richmond Agitation and Sedation Scale for delirium screening.** *J Hosp Med* 2012,  
17 **7**(5):450-453.
- 18 87. Han JH, Wilson A, Ely EW: **Delirium in the older emergency department patient:**  
19 **a quiet epidemic.** *Emerg Med Clin North Am* 2010, **28**(3):611-631.
- 20 88. Adamis D, Martin FC, Treloar A, Macdonald AJD: **Capacity, consent, and selection**  
21 **bias in a study of delirium.** *Journal of Medical Ethics* 2005, **31**(3):137-143.
- 22 89. Han JH, Wilson A, Vasilevskis EE, Shintani A, Schnelle JF, Dittus RS, Graves AJ,  
23 Storrow AB, Shuster J, Ely EW: **Diagnosing delirium in older emergency**  
24 **department patients: validity and reliability of the delirium triage screen and the**  
25 **brief confusion assessment method.** *Ann Emerg Med* 2013, **62**(5):457-465.
- 26 90. Han JH, Wilson A, Vasilevskis EE, Storrow AB, Shintani A, Schnelle J, Graves AJ,  
27 Dittus RS, Ely EW: **The Validation of the Emergency Department Delirium Triage**  
28 **Screen in Older Emergency Department Patients.** *Ann Emerg Med* 2012, **60**(4):S29.
- 29 91. Marcantonio ER, Ngo LH, O'Connor M, Jones RN, Crane PK, Metzger ED, Inouye  
30 SK: **3D-CAM: derivation and validation of a 3-minute diagnostic interview for**  
31 **CAM-defined delirium: a cross-sectional diagnostic test study.** *Annals of Internal*  
32 *Medicine* 2014, **161**(8):554-561.
- 33 92. **National Hip Fracture Database (NHFD), Annual Report 2018**  
34 [[https://www.nhfd.co.uk/files/2018ReportFiles/NHFD-2018-Annual-Report-](https://www.nhfd.co.uk/files/2018ReportFiles/NHFD-2018-Annual-Report-v101.pdf)  
35 [v101.pdf](https://www.nhfd.co.uk/files/2018ReportFiles/NHFD-2018-Annual-Report-v101.pdf)] Accessed 1 March 2019.
- 36 93. Lagarto L, Albuquerque E, Loureiro D, Vieira F, Esteves P, Neves S, Teixeira-  
37 Verissimo M, Cerejeira J: **Arousal changes and delirium in acute medically-ill male**

1        **older patients with and without dementia: a prospective study during**  
2        **hospitalization.** *Aging Ment Health* 2018:1-8.

3        94.     Han JH, Vasilevskis EE, Schnelle JF, Shintani A, Dittus RS, Wilson A, Ely EW: **The**  
4        4        **Diagnostic Performance of the Richmond Agitation Sedation Scale for Detecting**  
5        5        **Delirium in Older Emergency Department Patients.** *Academic Emergency Medicine*  
6        6        2015, **22**(7):878-882.

7        7        95.     Grossmann FF, Hasemann W, Kressig RW, Bingisser R, Nickel CH: **Performance of**  
8        8        **the modified Richmond Agitation Sedation Scale in identifying delirium in older**  
9        9        **ED patients.** *Am J Emerg Med* 2017, **35**(9):1324-1326.

**Figure 1. The 4 'A's Test (4AT)**



(label)

**Assessment test  
for delirium &  
cognitive impairment**

Patient name:

Date of birth:

Patient number:

Date:

Time:

Tester:

**CIRCLE**

**[1] ALERTNESS**

*This includes patients who may be markedly drowsy (eg. difficult to rouse and/or obviously sleepy during assessment) or agitated/hyperactive. Observe the patient. If asleep, attempt to wake with speech or gentle touch on shoulder. Ask the patient to state their name and address to assist rating.*

Normal (fully alert, but not agitated, throughout assessment)	0
Mild sleepiness for <10 seconds after waking, then normal	0
Clearly abnormal	4

**[2] AMT4**

*Age, date of birth, place (name of the hospital or building), current year.*

No mistakes	0
1 mistake	1
2 or more mistakes/untestable	2

**[3] ATTENTION**

*Ask the patient: "Please tell me the months of the year in backwards order, starting at December." To assist initial understanding one prompt of "what is the month before December?" is permitted.*

Months of the year backwards	Achieves 7 months or more correctly	0
	Starts but scores <7 months / refuses to start	1
	Untestable (cannot start because unwell, drowsy, inattentive)	2

**[4] ACUTE CHANGE OR FLUCTUATING COURSE**

*Evidence of significant change or fluctuation in: alertness, cognition, other mental function (eg. paranoia, hallucinations) arising over the last 2 weeks and still evident in last 24hrs*

No	0
Yes	4

**4 or above:** possible delirium +/- cognitive impairment  
**1-3:** possible cognitive impairment  
**0:** delirium or severe cognitive impairment unlikely (but delirium still possible if [4] information incomplete)


**4AT SCORE**

**GUIDANCE NOTES**

Version 1.2. Information and download: [www.the4AT.com](http://www.the4AT.com)


The 4AT is a screening instrument designed for rapid initial assessment of delirium and cognitive impairment. A score of 4 or more suggests delirium but is not diagnostic: more detailed assessment of mental status may be required to reach a diagnosis. A score of 1-3 suggests cognitive impairment and more detailed cognitive testing and informant history-taking are required. A score of 0 does not definitively exclude delirium or cognitive impairment: more detailed testing may be required depending on the clinical context. Items 1-3 are rated *solely on observation of the patient at the time of assessment*. Item 4 requires information from one or more source(s), eg. your own knowledge of the patient, other staff who know the patient (eg. ward nurses), GP letter, case notes, carers. The tester should take account of communication difficulties (hearing impairment, dysphasia, lack of common language) when carrying out the test and interpreting the score.

**Alertness:** Altered level of alertness is very likely to be delirium in general hospital settings. If the patient shows significant altered alertness during the bedside assessment, score 4 for this item. **AMT4 (Abbreviated Mental Test - 4):** This score can be extracted from items in the AMT10 if the latter is done immediately before. **Acute Change or Fluctuating Course:** Fluctuation can occur without delirium in some cases of dementia, but marked fluctuation usually indicates delirium. To help elicit any hallucinations and/or paranoid thoughts ask the patient questions such as, "Are you concerned about anything going on here?"; "Do you feel frightened by anything or anyone?"; "Have you been seeing or hearing anything unusual?"

**Figure 2. Study design overview flowchart**


**Figure 3.**

**STARD diagram of flow of participants through the study (total across all three sites)**





**Figure 4. Receiver Operator Characteristic Curve for 4AT diagnostic accuracy**



4AT scores range from 0-12. The cut-point of >3 is used in the scoring scheme to denote likely delirium. The 4AT scores are considered against the reference standard delirium assessment.





Click here to access/download  
**Supplementary Material**  
Additional Table 2.rtf







Click here to access/download  
**Supplementary Material**  
Additional Table 5.docx